

ACTELLIC[®] 25 EC

A wide spectrum organophosphate insecticide for use in controlling thrips, aphids, whiteflies, caterpillars, mites and weevils in agricultural, horticultural crops and insects in storage.

REGISTRATION NO. PCPB(R) 0015

GUARANTEE: 250 g/lit. pirimiphos-methyl

Storage: Store in a cool dry place away from sun and damp, out of reach of children, animals, food and feed.

Shelf life: At least two years from date of manufacturing, in original sealed containers.

FORMULATED in Kenya under license of Syngenta Agro AG, Basle, Switzerland

DISTRIBUTED IN KENYA BY:
TWIGA CHEMICAL INDUSTRIES LIMITED,
P.O. BOX 30172, NAIROBI, KENYA.
TEL: 338333

READ LABEL AND LEAFLET BEFORE USING THE PRODUCT


KEEP LOCKED OUT OF REACH OF CHILDREN

COMMERCIAL AND AGRICULTURAL CLASS

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

© = registered Trademark of a Syngenta Group Company

syngenta
Colour: YELLOW



ACTELLIC is a broad-spectrum insecticide with both contact and fumigant action for control of major insect pests in horticultural and agricultural crops and in stored grain. When used according to instructions no harmful residues will occur. Treated grain may be safely consumed by humans and livestock.

Directions for Use

To avoid risks to man and the environment comply with the use instructions.

Stored Grain – Bulk: Mix 20 ml of ACTELLIC in 25 litres water. Apply 1 litre per tonne of grain as it flows to the bulk store.

Bagged Grain: Mix 400 ml in every 20 litres of water and spray over 100 m² of surface (wet to run off). Spray also the ground surrounding the store. Ventilate store before re-entry.

NOTE: This formulation of ACTELLIC will not give control of the Larger Grain Borer (*Prostephanus truncates*). If infestation by this pest is suspected seek expert advice. ACTELLIC will give control of malathion resistant pests.

Potatoes/Vegetables including peas, beans – aphids, whitefly, thrips, spider mites, weevils, caterpillars, cutworms. Spray at first signs of pest infestation and repeat at 14-day spray intervals. Do not spray when leaves are wet or plants are stressed. Apply 2 lt./ha in 250-500 lt./ha (150 ml/20 lt.). Observe a pre-harvest interval of 7 days. For cutworm control apply as a band spray along the rows.

Apples, Citrus, Bananas, Pineapples and other fruits – caterpillars, scales and mealy bugs. Do not spray when leaves are wet or plants are stressed. Apply 3 lt./ha in 1000 lt./ha (60 ml/20 lt.). Observe a pre-harvest interval of 7 days.

Rice stem borer, Plant and leafhopper - Spray 2 lt. in 1000 litres of water per 1000m² (40 ml/20 lt., approximate coverage 6 x 6 paces).

Handling Precautions

Caution – irritating to eyes and skin. Wear rubber gloves, face shield and boots when handling the concentrate. Do not use if under medical advice

not to work with organophosphates. Do not eat, smoke or drink while working with the product; and before doing so wash hands and face. Avoid spray mist. Change clothes and wash body after work.

Storage

Store in original sealed containers away from children, food, feed and livestock. Do not store in stores with a temperature above 35°C. In case of fire extinguish with dry chemical, foam or water spray. Do not use water jet.

Environmental Hazards

Toxic to fish. Do not contaminate rivers, dams and ponds. Toxic to bees. Do not spray when bees are actively foraging.

First Aid

Eye splashes: Wash eyes with plenty of clean water or eyewash for 10-15 minutes. See a doctor. **Skin splashes:** wash off immediately with water and soap. Remove contaminated clothing. If swallowed, do not make the patient vomit. Remove patient immediately from contaminated area and keep the patient in a cool and comfortable place. Contact a doctor immediately

Toxicological information

If poisoning is suspected, Atropine should be administered immediately at 5 mg intramuscularly, then 2 mg at 30 minutes intervals. At the same time Fralidoxine (p-25 or p-2m) should be given, 1 g intramuscularly in 2-3 ml water. No sedative should be given. Repeat doses depending on clinical response.

Symptoms of poisoning

These include nausea, blurred vision, tightness in chest, headaches, loss of co-ordination and abdominal cramp. Severe poisoning may lead to respiratory arrest. Oxygen and artificial respiration may be required.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

User's Risk

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Before using the product, please read the attached product label and accompanying packaging leaflet.

Use pesticides safely.

© = registered trademark of Syngenta Crop Protection AG, Basle,

Emergency Call:

Syngenta Alarm Centre, Huddersfield, UK
Phone: + 44 1484 538 444
Fax: + 44 1484 433 197