



Bulldock[®]

New generation Bayer pyrethroid

The success story continues

Broad spectrum insecticide
for the control of biting and sucking insects.



Bayer



Expertise with Responsibility

Bulldock[®] 025 EC

Beta-cyfluthrin 25gms per litre

Bulldock[®] 025EC is a highly effective broad spectrum contact as well as stomach poison insecticide in the class of modern pyrethroid.

Main Features

Bulldock[®] is characterised by a very broad spectrum of activity for controlling leaf-eating

- ▶ caterpillars,
- ▶ bollworms,
- ▶ beetles and their larvae



Coffee Leafminer




Cabbage moth


and sucking insects such as

- ▶ aphids,
- ▶ leafhoppers,
- ▶ thrips

Bulldock[®] is used in all important crops such as Cotton, Vegetables, Ornamentals, Coffee and other crops.



Diamondback moth



Bollworm

Advantages of using Bulldock

Rate of use

Higher concentration of active ingredients hence, less dosage.

Mode of action

- ▶ Excellent effect on biting and sucking insects.
- ▶ Act as a contact as well as a stomach poison.
- ▶ Combines fast initial knock down with long-lasting residual activity for 14 days.
- ▶ Bulldock is harmful to bees when in contact with the spray



Aphid

The success story continues



Bulldock® 025 EC

Beta-cyfluthrin 25gms per litre



Cotton stainer

Outstanding performance

Provides excellent control of pests that are less well controlled by other products (e.g. the cotton boll weevil).

Crop tolerance

Well tolerated by plants when used at recommended rates.

Compatibility


Excellent mixing compatibility with Bayer insecticides and fungicides e.g. Antracol, XenTari, Milraz, Bayleton, Folimat, Metasystox, Bayfolan, e.t.c.

Application Rates and Timing

Crop	Pests	Rate lt/Ha	Rate per 20 lts Water	Timing
Coffee	Leafminer	0.5	10 ml	Start spraying when the insects are seen and repeat as required.
Vegetables	African bollworm, Diamond backmoth, Aphids, Cabbage Moth.	0.6	15 ml	Start spraying when the insects are seen and ,repeat as required. For effective control of Diamondback moth alternate or mix with XenTari BT.
Cotton	African bollworm, Pink bollworm, Stainer bugs, Aphids, Whiteflies, Leafworms, Lygus spp.	0.6	15 ml	Start spraying when the insects are seen and repeat as required.
Ornamentals	Catepillars, African Bollworm, Budfly.	0.6	16 ml	Start spraying when the insects are seen and repeat as required

Water volume : 1,000 litres per hectare.

PHI: Vegetables 3 days



Bulldock® 025 EC

Beta-cyfluthrin 25gms per litre



**Bulldock® is a registered trademark
of Bayer AG, Germany.**

Enquiries:

**BAYER EAST AFRICA LTD.
BUSINESS GROUP CROP PROTECTION**
P.O. Box 30321, 00100-Nairobi, Kenya.

HEAD OFFICE:

Outering Road, Ruaraka
Tel: 860667 - 74, Fax: 803345 / 861836.

FACTORY:

Lunga Lunga Road (Industrial Area)
Tel: 531272 - 5, Fax: 544917.

Email: crop@bayerea.com

PRECAUTIONS

**Users of this product are reminded to always
refer to the product label for guidance on proper
use and handling of the pesticide.**