

## DIRECTIONS FOR USE

Measuring cups are provided with **FUNGURAN-OH** to enable the user to measure the correct amount of product into a 20 Litre Debe or Knapsack Sprayer.

When preparing the spray mixture it is advisable to mix the required amount of **FUNGURAN-OH** into a smooth thin paste with a small quantity of water in the bottom of the Debe. Then add the correct quantity of water and stir well before pouring into the Knapsack Sprayer.

Ensure even spray coverage of the target crop. All surfaces of the foliage must be covered to prevent disease developing.

## APPLICATION RATE AND TIMING OF APPLICATION

CROP	DISEASE	TIMING	RATES OF USE	
			KG/HA	GMS PER 20 L WATER
COFFEE	LEAF RUST	before onset of rains and at 21-28 day intervals during rains. (Normally 2 applications during short rains and 5 applications during long rains will be necessary.)	3.5 in 1000 litres water/ha	70 1 CUP
	COFFEE BERRY DISEASE	1st spray after flowering, before onset of rains. Thereafter spray at 21-28 day intervals until picking time.	7.0 in 1000 litres water/ha	140 2 CUPS
BEANS	BACTERIAL BLIGHT	Spray when disease anticipated. Repeat at 10-14 day intervals.	1-2	40-80 1 CUP
TOMATOES	EARLY BLIGHT	Spray at 7-14 day intervals during infection period.	1-2	40-80 1 CUP
POTATOES	EARLY BLIGHT	Spray at 7-14 day intervals during infection period.	1-2	50 1 CUP
GRAPES	DOWNY MILDEW	Spray at 7-14 day intervals during infection period.	1-2	50 1 CUP
CITRUS	BLACK SPOT	Spray at 75% Petal Drop. Repeat at 28 day intervals.	2.0	40 1 CUP
ORNAMENTALS	DOWNY MILDEW	Apply before disease appears. Repeat at 10-14 day intervals. Can be tank mixed with specific fungicides or used as an alternative fungicide in a spray programme.	2-4	50-100 1 CUP

## COMPATIBILITY

**FUNGURAN-OH** is compatible with many fungicides and insecticides e.g. FENITROTHION, CHLOROTHALONIL, PROPICONAZOLE, DITHIANON, DELTAMETHRIN.

## PRECAUTIONS

Do not inhale powder or spray mist. Avoid contact with skin or eyes. Do not swallow. Prevent spray drift on to other crops, areas not under treatment or into rivers or dams.

## FIRST AID IN CASE OF POISONING

Wash skin with soap and water and wash eyes copiously with water.

## STORAGE INSTRUCTIONS

Store under dry conditions and avoid excessive heat.

## TOXICITY

**FUNGURAN-OH** is defined as a toxicity class III product in the WHO classification.

## JINSI YA KUTUMIA

Vipimo vimo ndani kumwezesha mtumiaji kupima dawa sawa sawa kwenye. Debe la lita ishirini au kifaa cha kupulizia dawa Wakati wa kutayarisha mchanganyiko wa dawa, imependekeza mtumiaji achanganye kiwango kinachohitajika na maji kidogo hadi mchanganyiko uwe laini kwenye debe halafu aongeze maji kabla ya kutia kwenye kifaa cha kupulizia dawa. Hakikisha dawa imetapakaa sawa sawa katika mimea ili kuzuia ugonjwa.

## KIWANGO NA WAKATI WA MATUMIZI

MIMEA	UGONJWA	WAKATI	KIWANGO KG/HA	CHA MATUMIZI GMS/20 L MAJI
KAHAWA	KUTU YA MAJANI	Kabla ya mvua kuanza kisha baada ya siku 21-28 wakati wa mvua (kawaida tumia mara mbili wakati wa mvua ndogo na mara tano wakati wa mvua za masika)	3.5 kwa 1000 L	70 KIKOMBE 1
	UGONJWA WA MBEGO ZA KAHAWA (UKUNGU)	Tia dawa baada ya maua kuchanua, kabla ya mvua kuanza halafu kila baada ya siku 21-28 hadi wakati wa wakati wa kuvuna	7.0 kwa 1000 L	140 VIKOMBE 2
JAMII YA MAHA-RAGWE	KUTU KUGA OR KUTU KUVU	Nyunyiza wakati unatazamia ugonjwa kuanza kisha kila baada ya siku 10-14	1-2	40-80 KIKOMBE 1
NYANYA	KUTU KUGA OR KUTU KUVU	Nyunyiza kila baada ya siku 7-14 wakati wa kuambukizwa	1-2	40-80 KIKOMBE 1
VIAZI	KUTU KUGA	Nyunyiza kila baada ya siku 7-14 wakati wa kuambukizwa	1-2	40-80 KIKOMBE 1
ZABIBU	KUVU YA UNGA	Nyunyiza kila baada ya siku 7-14 wakati wa kuambukizwa	1-2	50 KIKOMBE 1
MUCHUNGWA	BLACK SPOT	Nyunyiza kwenye majani kwa 75% kisha kila baada ya siku 20-28	2.0	40 KIKOMBE 1
MAUJA YA	KUVU YA UNGA	Nyunyiza kabla ya kuambukizwa kisha kila baada ya siku 10-14. Ina weza kuchanganywa na dawa fulani nyingine kwa birika la maji au kutumiwa kwenye kifaa cha kupulizia dawa.	2-4	50-100 KIKOMBE 1

## UTANGAMANJI

**FUNGURAN-OH** inaweza kuchanganywa n adawa nyingi za kuuwa wadudu na viini kama: FENITROTHION, CHLOROTHALONIL, PROPICONAZOLE, DITHIANON, DELTAMETHRIN.

## KINGA

Usinuse dawa. Jikinge isikugewe ngozi wala macho. Pis Jihadhari na kumeza dawa. Hakikisha dawa isienee kwenye mimea mengine, eneo lililo hitaji kunyunyiziwa dawa na kwenye mito au eneo lolote la maji.

## HUDUMA YA KWANZA IWAPO MTU ATAPATA SUMU

Osha ngozi kwa sabuni na maji. Halafu macho yatie maji kwa wingi.

## JINSI YA KUHIFADHI DAWA

Hifadhi dawa pahali pakavu pasipo na joto jingi.

## KIWANGO CHA SUMU

**FUNGURAN-OH** iko kwenye kiwango cha daraja ya tatu kwenye mpangilio wa WHO (Idaro ya Afya ya Umoja wa Mataifa).

## WARRANTY STATEMENT

SPIESS-URANIA CHEMICALS GMBH warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of SPIESS-URANIA CHEMICALS GMBH. In no case shall SPIESS-URANIA CHEMICALS GMBH be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks are assumed by the buyer. SPIESS-URANIA CHEMICALS GMBH MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Shelf Life: 5 years